EFFECT OF HEALTH LITERACY ON CHRONIC HEPATITIS B VIRUS INFECTION MANAGEMENT AND PROGNOSIS

Ceren YILMAZ ^[0], Fehmi ATES ^[0], Hakan DEMIRCI ^[0]

¹ Department of Internal Medicine, Mersin University Medical Faculty, Mersin, TURKEY

² Department of Gastroenterology, Mersin University Medical Faculty, Mersin, TURKEY

² Department of Family Medicine, University of Health Sciences Turkey Bursa Yuksek Ihtisas Training and Research Hospital, Bursa, TURKEY

42

ABSTRACT

Aim: In this study, we aimed to examine the relationship of health literacy with the diagnostic stage of the disease, the risk of encountering complications, and other prognostic variables in chronic hepatitis B patients.

Methods: A total of 264 patients (146 male and 118 female) with chronic hepatitis B disease who presented to the gastroenterology outpatient clinic of Mersin University Faculty of Medicine Hospital in 2019 were included in the study. The HLS-EU-47 scale was used to measure the level of health literacy.

Results: The health literacy of patients taking antivirals was significantly lower than those who did not use them (p=0.031). The health literacy of patients who underwent regular follow-up after chronic hepatitis B was significantly higher than those who did not (p<0.001). The health literacy of patients with the chronic form of hepatitis B was significantly higher than the others (p<0.001). The health literacy of patients with cirrhosis development was significantly lower than those without (p<0.001). In the group that needed paracentesis, health literacy was observed to be lower than in the group that did not need paracentesis (p<0.004). The health literacy of patients hospitalized in the last year was significantly lower than that of non-hospitalized patients (p<0.002). The number of hospitalizations in the last year was associated with health literacy (p<0.009).

Conclusions: Health literacy is associated with the course of the disease in chronic hepatitis B patients. Frequent hospitalization, frequent visits to the emergency department, the need for frequent paracentesis, and the need for antiviral therapy were more common in the group with low health literacy. Improving health literacy will positively affect the prognosis of these patients.

Keywords: hospitalization; chronic hepatitis B; paracentesis; prognosis; health literacy

Corresponding Author: Ceren YILMAZ ylmz_ceren@hotmail.com

Received: December 25, 2023; Accepted: December 29, 2023; Published Online: December 30, 2023

Cite this article as: Yilmaz, C., Ates, F. & Demirci, H. (2023). Effect of Health Literacy on Chronic Hepatitis B Virus Infection Management and Prognosis. European Health Literacy Journal 3(4), 42-50.



INTRODUCTION

The World Health Organization (WHO) estimates that 296 million people were living with chronic hepatitis B infection in 2019, with 1.5 million new infections added every year. It is estimated that 820,000 deaths occurred because of hepatitis B in 2019, mostly due to cirrhosis and hepatocellular carcinoma. As of 2019, worldwide awareness of hepatitis B was 10.5%, while 22% of the people diagnosed were receiving treatment [1].

The progression of cirrhosis can be slowed down, the incidence of liver cancer can be reduced, and improvement in long-term survival can be achieved with the treatment of chronic hepatitis B infection. The WHO has estimated that 12% to 25% of people with chronic hepatitis B infection will require antiviral treatment in 2021, depending on the settings and eligibility criteria. Guidelines recommend the use of oral medications (tenofovir or entecavir) as the most potent treatment modality to suppress the hepatitis B virus. Many of the patients who start antiviral treatment for hepatitis B virus must continuethetreatment for life [2].

Most people diagnosed with liver cancerin low-income settings die within a short time. In high-income countries, patients with symptoms present to the hospital earlier, and have easier access to surgery and chemotherapy, which can prolong life for months to years. Liver transplantation is sometimes performed for people with cirrhosis or liver cancer in high-income countries.³ Health literacy can be defined as the capacity to acquire, understand, and interpret basic health information and services in order to protect and improve human health and improve deteriorating Health [2]. Health literacy includes having the ability to make decisions and being able to use this ability in health-related issues [3-5]. Those with low health literacy have difficulty understanding health information and following the necessary instructions, and have problems getting adequate health care [6-10].

Among the chronic diseases, health literacy is most commonly investigated in association withdiabetes mellitus. chronic obstructive pulmonary disease, obesity, and hypertension. It has been reported that people with better health literacy as regards these chronic diseases are easier to treat. face fewer complications, or are diagnosed earlier [11-14]. A limited number of studies on the relationship of health literacy with chronic hepatitis B disease and its complicationsare present in the literature [15-19].

The aim of the current study was to evaluate the relationship of health literacy with the diagnostic stage of the disease, the risk of encountering complications, and other prognostic variables in chronic hepatitis B patients.

METHODS

Approval for the study was obtained from the hospital's ethics committee. A total of 146 male and 118 female patients with chronic hepatitis B disease who presented to the gastroenterology outpatient clinic of Mersin University Faculty of Medicine Hospital in 2019 were included in the study. A questionnaire on the sociodemographic characteristics of the patients used. and the HLS-EU-47 scale was wasemployed to measure the level of health literacy [20]. The questionnaires were applied with the face-to-face interview technique. Patients who could not communicate verbally, did not have the mental capacity to understand and respond to the questions of the European Health Literacy Survey Questionnaire 47 (HLS-EU-Q47), or did not agree to participate in the study were excluded from the study.

Diagnostic criteria for chronic hepatitis B

These were as follows:

- HBsAg positive in the blood for more than 6 months,

- Serum HBV DNA>20,000 IU/ml in HBeAg-positive patients,

- Serum HBV DNA >2,000 IU/ml in HBeAg-negative patients,

- Persistent or intermittent high levels of AST and ALT,

- A liver biopsy showing signs of chronic hepatitis B disease with moderate or severe necroinflammation.

HLS-EU-47 Scale

The HLS-EU form consists of 47 questions. Each of the 47 questions was evaluated on a scale scored between 1 and 4 (1: very difficult, 2: difficult, 3: easy, 4: very easy). This scale is based on the assessment of how difficult or easy the behavior specified in each question is according to one's own perception. Within the framework of the HLS-EU-Q47 concept, sub-indexes are created by taking the average values of the health literacy parameters [20].

Statistical Analysis

The data obtained were analyzed Sciences (SPSS). With all statistical analyses, p<0.05 was accepted as the significance level and relationships were evaluated within the 95% confidence interval. Descriptive findings of the werenumber and study group percentage distributions for categorical variables, and mean, standard deviation, and medianfor continuous variables. The percentilesare given by their lowest and highest values. The relationships between the numerically dependent and independent variables were analyzed by Spearman correlation analysis. Shapiro-Wilk test and Kolmogorov-Smirnov test were used to assess whether the variables followed a normal distribution. According to the normality test results, the Mann-Whitney U and Kruskal Wallis tests were used to compare the groups.

RESULTS

The comparisons between the general characteristics of the participants and the HLS-EU-Q47 results are given in Table 1. Accordingly, the health literacy of university and high school graduates was found to be significantly higher than those who were illiterate oronly had primary education. The health literacy of working participants was found to be significantly higher than that of non-working participants. The health

literacy of public employees was found to be higher than that of housewives and the selfemployed groups. It was observed that the health literacy of the participants with good socioeconomic status was higher than those withalow socioeconomic status. The health literacy of those with a high body mass index was significantly lower compared to those with a low body mass index. The health literacy of older participants was significantly lower thanyounger participants.

		European Health Literacy Scale Scores					
		Mean	Standard Deviation	Median	Minimum	Maximum	p-value
Condon	Male	30.48	9.18	30.14	10.64	46.81	0 2028
Gender	Female	29.22	9.17	29.26	9.22	46.81	
	Single	29.82	9.17	30.65	12.77	46.10	
Marital Status Education Occupation	Married	30.12	9.07	30.14	9.22	46.81	0.316 ^b
	Divorced	25.30	11.40	21.19	10.64	44.68	
Education Occupation	Illiterate	19.09	5.97	17.91	9.22	31.56	
	Primary	25.35	7.14	24.11	10.64	42.91	-0.001b
	High School	32.86	8.05	34.04	13.48	46.81	<0.001
	University	38.51	5.00	38.83	25.53	46.81	
	Yes	31.94	9.20	34.04	10.64	46.81	-0.001 ⁸
Occupation	No	27.98	8.75	27.31	9.22	46.81	<0.001
	Self-employed	28.68	9.13	28.19	10.64	46.10	
	Officer	36.91	5.93	37.59	14.54	45.39	
Occupation	Retired	31.40	8.80	32.26	14.54	46.81	<0.001 ^b
	Housewife	25.50	8.45	24.47	9.22	44.68	
	500-1000 ₺	20.12	6.47	17.73	12.77	30.14	
 	1001-2000 赴	25.62	7.75	24.65	10.64	45.74	<0.001 ^b
Income	2001-3000 赴	32.56	8.46	34.40	12.77	46.81	<0.001
	>3000 ₺	37.05	6.33	37.59	19.50	46.81	
	Yes	29.65	8.73	29.43	10.64	46.81	
Smoking	No	30.13	9.45	30.85	9.22	46.81	0.869^{b}
	Quit	29.39	9.17	27.66	13.48	45.39	
	Yes	34.26	10.14	36.88	16.31	45.04	
Alcohol	No	29.96	9.17	30.32	9.22	46.81	0.127 ^b
	Quit	26.09	7.77	25.00	15.96	41.49	
Chronic	Yes	29.26	9.27	28.55	9.22	46.81	
Disease (Other than Chronic Hepatitis)	No	31.20	8.89	34.04	10.64	46.10	0.100^{a}
		Correlat	tion Coefficien	nt (Rho)			
Age		-0.166					0.007
Body Mass Inc	dex (kg/m^2)	-0.195					0.001

^a: Mann Whitney u test; ^b: Kruskal Wallis H test; rho = Spearman Correlation Coefficient

The mean score of the participants on the health literacy scale was calculated as 29.92±9.18. Table 2 presents the comparisons of the health literacy scores with hepatitis B variables and Table 3 presents the comparisons of the health literacy scores with cirrhosis complications. The health literacy of patients taking antivirals was significantly lower than of those who were not use them. The health literacy of patients who underwent regular follow-up after chronic hepatitis B was significantly higher than those who did not. The health literacy of patients continuing in the form of chronic chronic hepatitis B was significantly higher than those who did not. The health literacy of patients with the cirrhosisdevelopment was significantly lower than those without this complication. The relationship between paracentesis needs and health literacy of cirrhosis patients with ascites is shown in Figure 3. In the group that needed paracentesis, health literacy was observed to be lower than in the group that did not need paracentesis.

Table 2. Comparisons of the health literacy scores with variables related to Hepatitis B

		Mean	Standard Deviation	Median	Minimum	Maximum	p-value
Using	Yes	29.35	9.21	28.72	10.64	46.81	0.0218
Using Antiviral drug M Regular M follow-up M Persisting as M Chronic M Hepatitis B Cirrhosis M Cirrhosis C Stage M HCC M	No	32.50	8.66	34.04	9.22	46.81	-0.031
Regular	Yes	31.56	8.68	32.44	10.64	46.81	<0.0018
follow-up	No	24.98	8.91	24.64	9.22	42.91	<0.001
Persisting as	Yes	32.70	9.08	34.04	14.18	46.81	
Chronic Hepatitis B	No	26.15	8.60	26.06	9.22	45.74	<0.001 ^a
follow-up Persisting as Chronic Hepatitis B Cirrhosis Cirrhosis	Yes	26.26	8.66	26.06	9.22	45.74	<0.0018
	No	32.70	8.58	34.04	14.18	46.81	<0.001
Cirrhosis	Compensated	26.75	8.98	28.01	9.22	45.74	0.2618
Stage	Decompensated	24.26	6.68	30.14	12.77	35.82	-0.201
	Yes	22.70	9.17	17.73	14.54	35.82	
НСС	No	30.06	9.14	30.14	9.22	46.81	0.075 ^a

^a: Mann Whitney u test; HCC: Hepatocellular carcinoma

The comparisons between health literacy scores and hospitalization history is given in Table 4. The health literacy of patients hospitalized in the last year was significantly lower than that of non-hospitalized patients. The number of hospitalizations in the last year also decreased as health literacy increased.

		Mean	Standard Deviation	Median	Minimum	Maximum	p-value
Anitan	No	29.81	9.22	30.14	9.22	46.81	0.425 ^a
Ascilles	Yes	31.60	8.48	32.09	16.67	41.49	-0.423
Variana	No	29.79	9.07	30.14	9.22	46.81	0.506 ^a
varices	Yes	31.07	10.23	30.83	14.54	45.74	-0.300
Varices	No	29.86	9.18	30.14	9.22	46.81	0.6118
hemorrhage	Yes	31.21	9.51	35.82	16.67	41.84	-0.011
Domitomitic	No	29.82	9.18	30.14	9.22	46.81	
Peritonius	Yes	-	-	-	-	-	7-
En e en la el en et less	No	29.89	9.10	30.14	9.22	46.81	0.6008
Encephalopathy	Yes	31.03	13.14	35.46	10.64	42.20	-0.009
Hospitalization due	Yes	22.55	7.43	23.40	14.54	33.33	0.066 ^a
to encephalopathy	No	30.06	9.16	30.14	9.22	46.81	-0.000
	Yes	21.32	5.31	21.28	14.54	28.72	0.004a
Paracentesis	No	30.22	9.14	30.85	9.22	46.81	-0.004
Rate of paracentesis	>1 in a week	20.92	9.64	20.92	12.77	31.56	
	Once in a week	20.28	6.14	16.67	14.54	28.72	0.724 ^b
	Once in a month	22.61	4.58	21.99	17.13	28.72	
	No	-	-	-	-	-	

Table 3. Comparisons of the health literacy scores with cirrhosis complications

^a: Mann Whitney u test; ^b: Kruskal Wallis H test

Table 4. Comparisons between health Literacy scores and hospitalization history

		Mean	Standard Deviation	Median	Minimum	Maximum	p-value
Hospitalization	No	30.87	9.16	31.56	10.64	46.81	0.0028
within the last year	Yes	26.76	8.56	25.72	9.22	45.04	-0.002
	None	30.87	9.16	31.56	10.64	46.81	
	Once	26.76	8.98	27.31	9.22	45.04	
within the last year	More than once	26.77	7.52	25.18	16.67	42.91	0.009 ^b
	None	30.68	9.28	32.09	10.64	46.81	
Visits to the emergency department within the last year	Once	29.41	9.38	28.72	13.48	44.33	
	More than once	28.12	7.99	27.31	9.22	45.74	0.210 ^b

^a: Mann Whitney u test; ^b: Kruskal Wallis H test

DISCUSSION

The health literacy of patients taking antivirals was significantly lower than those who did not use them. The health literacy of patients who underwent regular follow-up after chronic hepatitis B was significantly higher than those who did not. Also, the health literacy of patients continuing in the form of chronic hepatitis B was significantly higher than those who did not. In addition, the health literacy of patients with the development of cirrhosis was significantly lower than that those without his complication. In the group that needed paracentesis, health literacy was observed to be lower than in the group that did not need paracentesis. The health literacy of patients hospitalized in the last year was significantly lower than that of non-hospitalized patients. The number of hospitalizations in the last year was similarly related to health literacy.

In previous studies, it was found that seeking treatment and the adherence to medication were high in individuals with good health literacy for chronic diseases [9, 13, 21, 22]. In this study, antiviral use was associated with low health literacy. The guidelines recommend antiviral therapy and close monitoring of patients in cases of active chronic hepatitis B. Since there are patients who are admitted to a university hospital, they can be considered patients who are looking for treatment. However. the patients recommended antiviral treatment have slightly worse progression and are probably cases that have remained away from regular follow-up.

In one study, the researchers found that among untreated patients, the annual cirrhosis and HCC incidence ranged from 0.26% to 1.30% and 0.04% to 3.80% in inactive patients, and 0.55 to 4.05% and 0.19 to 6.03% in active patients, respectively [23]. They claimed that transition rates increased with age, male sex, the presence of fibrosis/cirrhosis, and active disease and/or antiviral treatment. It is known that these transitions are less common in cases that have regular follow-up and comply with the recommendations. Although unique the characteristics of the virus and the host also affect the transition to cancer or cirrhosis, the transition to advanced stage disease slows down in individuals with good health literacy. This situation can be interpreted in two ways. First, individuals with low health literacy should be monitored more closely. Secondly, the prognosis and survival of patients can be improved with health literacy training.

It is also expected that the need for paracentesis will be lower in cases with good health literacy, which is a positive result. Patients can achieve a decrease in the need for paracentesis by following the warnings about nutrition and taking medications. In the studies conducted, the need for paracentesis was associated with multiple factors [24]. One of the most important results of the research is that paracentesis, which is an interventional practice and indicates a poor prognosis, is associated with health literacy. In addition, paracentesis is also associated with concomitant cardiac problems, and health literacy has also been known to be protective in patients with heart disease in previous studies.

There is a lot of research available on hospitalization and health literacy [25-27]. Frequent hospitalization of patients is a sign that the course of the disease is poor. Disease severity has been associated with an increase in the frequency of hospitalization in patients. As a result of the research, hospitalization, which is one of the signs of a poor prognosis, was found to be associated with health literacy. Hospitalization is a sign that the course of the disease is bad, and it is also accompanied by an increase in medical expenses. Considering the costs of hospitalization, paracentesis, and antiviral treatment, it can be predicted that health literacy will save both the individuals and governments from a significant financial burden.

Limitations

Our research was conducted in a singlecenter manner, and generalizations cannot be made. The number of our cases remained low due to the Covid pandemic. The limited number of participants also limited the study. It may be a bias to conduct research in a third-line university hospital. The health literacy of patients who came to the university hospital may be different from the patients who did not care about the treatment at all or who had low treatment compliance. For this reason, our collection of cases at the university hospital outpatient clinic may have negatively affected our results.

Conclusions

Health literacy is associated with the course of the disease in chronic hepatitis B patients. Frequent hospitalization, frequent visits to the emergency department, the need for frequent paracentesis, and the need for antiviral therapy are observed more commonly in the group with low health literacy. Improving health literacy will positively affect the prognosis of these patients. It may be necessary to be more descriptive in patients with poor health literacy. Patients with chronic hepatitis B who have been found to have poor health literacy should be monitored more closely, and support from their companions could be requested for their treatment and control.

Conflicts of interest

The authors have nothing to disclose.

Funding

Nothing to declare.

REFERENCES

1. World Health Organization. Hepatitis B. 2022. Access Date: 28.12.2023. Available from: https://www.who.int/news-room/fact-

sheets/detail/hepatitis-b.

2. Liu, C., Wang, D., Liu, C., Jiang, J., Wang, X., Chen, H., ... & Zhang, X. (2020). What is the meaning of health literacy? A systematic review and qualitative synthesis. *Family medicine and community health*, 8(2).

3. Corrarino, J. E. (2013). Health literacy and women's health: challenges and opportunities. *Journal of midwifery & women's health*, 58(3), 257-264.

4. Speros, C. (2005). Health literacy: concept analysis. *Journal of advanced nursing*, *50*(6), 633-640.

5. Berkman, N. D., Davis, T. C., & McCormack, L. (2010). Health literacy: what is it?. *Journal of health communication*, 15(S2), 9-19.

6. Chew, L. D., Bradley, K. A., & Boyko, E. J. (2004). Brief questions to identify patients with inadequate health literacy.

7. McCray, A. T. (2005). Promoting health literacy. *Journal of the American Medical Informatics Association*, *12*(2), 152-163.

8. Bozkurt, H., & Demirci, H. (2019). Health literacy among older persons in Turkey. *The Aging Male*, 22(4), 272-277.

9. Cayci, H. M., Erdogdu, U. E., Demirci, H., Ardic, A., Topak, N. Y., & Taymur, İ. (2018). Effect of health literacy on help-seeking behavior in morbidly obese patients agreeing to bariatric surgery. *Obesity surgery*, 28, 791-797.

10. Azkan Ture, D., Bhattacharya, S., Demirci, H., & Yildiz, T. (2022). Health literacy and health outcomes in chronic obstructive pulmonary disease patients: an explorative study. *Frontiers in Public Health*, *10*, 846768.

11. Guclu, O. A., Demirci, H., Ocakoglu, G., Guclu, Y., Uzaslan, E., & Karadag, M. (2019). Relationship of pneumococcal and influenza vaccination frequency with health literacy in the rural population in Turkey. *Vaccine*, *37*(44), 6617-6623.

12. Ciftci, F., Demirci, H., Ciftci, H. N., & OCAKOĞLU, G. (2021). Validation of Turkish version of newest vital sign scale to assess health literacy. *Bezmialem Science*, 9(2), 219.

13. Biyik, I., Mut, A. N. U., Albayrak, M., Kucuk, B., Koras, O., Keskin, F., & Demirci, H. (2021). Effect of health literacy on help-seeking behavior: a comparison of patients accepting surgery and refusing surgery for urinary incontinence. *Journal of Gynecology Obstetrics and Human Reproduction*, *50*(4), 101908.

14. Erdoğdu, U. E., Çaycı, H. M., Tardu, A., Arslan, U., Demirci, H., & Yıldırım, Ç. (2020). Relationship between health literacy and quality of colonoscopy bowel preparation. *The Turkish Journal of Gastroenterology*, *31*(11), 799.

15. Dahl, T. F., Cowie, B. C., Biggs, B. A., Leder, K., MacLachlan, J. H., & Marshall, C. (2014). Health literacy

in patients with chronic hepatitis B attending a tertiary hospital in Melbourne: a questionnaire based survey. *BMC infectious diseases*, 14(1), 1-9.

16. Gulati, R., Nawaz, M., & Pyrsopoulos, N. T. (2018). Health literacy and liver disease. *Clinical Liver Disease*, *11*(2), 48.

17. Choudhry, F. R., Ming, L. C., Munawar, K., Zaidi, S. T. R., Patel, R. P., Khan, T. M., & Elmer, S. (2019). Health literacy studies conducted in Australia: a scoping review. *International journal of environmental research and public health*, *16*(7), 1112.

18. Castro-Sánchez, E., Chang, P. W., Vila-Candel, R., Escobedo, A. A., & Holmes, A. H. (2016). Health literacy and infectious diseases: why does it matter?. *International Journal of Infectious Diseases*, 43, 103-110.

19. Hajarizadeh, B., Wallace, J., Richmond, J., Ngo, N., & Enright, C. (2015). Hepatitis B knowledge and associated factors among people with chronic hepatitis B. *Australian and New Zealand Journal of Public Health*, *39*(6), 563-568.

20. Sørensen, K., Pelikan, J. M., Röthlin, F., Ganahl, K., Slonska, Z., Doyle, G., ... & Brand, H. (2015). Health literacy in Europe: comparative results of the European health literacy survey (HLS-EU). *The European journal of public health*, *25*(6), 1053-1058.

21. Ocakoglu, G., Demirci, H., GUCLU, O. A., & Guclu, Y. (2020). Association between health literacy and medication adherence in the elderly population with chronic disease. *Ethiopian Journal of Health Development*, 34(2).

22. Turkoglu, A. R., Demirci, H., Coban, S., Guzelsoy, M., Toprak, E., Aydos, M. M., ... & Ustundag, Y. (2018). Evaluation of the relationship between compliance with the follow-up and treatment protocol and health literacy in bladder tumor patients. *The Aging Male*.

23. Liu, M., Tseng, T. C., Jun, D. W., Yeh, M. L., Trinh, H., Wong, G. L., ... & Nguyen, M. H. (2021). Transition rates to cirrhosis and liver cancer by age, gender, disease and treatment status in Asian chronic hepatitis B patients. *Hepatology international*, *15*, 71-81.

24. Kalo, E., Sheriff, N., Isaac, M., Baig, A., Read, S., & Ahlenstiel, G. (2022). A Call for Implementation of an Evidence-Based, Quality Improvement, Decompensated Cirrhosis Discharge Care Bundle in Australia. *Livers*, *2*(2), 97-104.

25. Cartwright, L. A., Dumenci, L., Cassel, J. B., Thomson, M. D., & Matsuyama, R. K. (2017). Health literacy is an independent predictor of cancer patients' hospitalizations. *HLRP: Health Literacy Research and Practice*, 1(4), e153-e162.

26. Rahman, T. T. A. (2014). Health literacy: prevalence among elderly care givers and its impact on the frequency of elderly hospitalization and elderly health related quality of life. *Advances in Aging Research*, *3*(05), 380.

27. Gulati, R., Nawaz, M., & Pyrsopoulos, N. T. (2018). Health literacy and liver disease. *Clinical Liver Disease*, *11*(2), 48.